Emerging Innovations
SCIENCE, TECHNOLOGY & SOCIAL STUDIES

Frontiers in Social Thoughts and Humanity (ISSN: - 2978-3755)

Characteristics of Preparing Muay Thai Athletes for the Professional Competitive Level

Suleyman Suleymanov

Professional Muay Thai and MMA coach Niles, Illinois

Received: 15 April 2026
Revised: 15 May 2026
Accepted: 22 May 2026
Published: 09 June2026

Volume03 **Issue** 06, 2026
Page No - 01-11

Article Citation: Suleyman Suleymanov. (2026). Characteristics of Preparing Muay Thai Athletes for the Professional Competitive Level. Frontiers in Social Thoughts and Humanity, 3(06), 1–11. Retrieved from https://irjernet.com/index.php/fsth/article/view/581

Copyright: © 2026 The Authors. Published by FSTH under the Creative Commons CC BY License

Abstract

The international growth of professional Muay Thai has outpaced the scientific evidence available to coaches who prepare athletes for sanctioned five-round contests at elite venues such as Lumpinee, Rajadamnern and ONE Championship cards. This article aims to clarify the contemporary characteristics of preparation for the professional competitive level by synthesizing peer-reviewed evidence on physiological demands, periodized strength and conditioning, technical-tactical pacing, weight management, injury epidemiology and recovery monitoring. The methodology is a structured narrative review combined with comparative analysis of twenty primary sources published from 2020 onward and indexed in Scopus, PubMed and Web of Science, supplemented by two industry market reports. Results indicate that elite preparation rests on four interacting domains: physical capacity, technical and tactical execution, body composition with periodized nutrition, and recovery monitoring through heart rate variability. Recent intervention data show that sport-specific high-intensity interval training improves Wingate peak power by about 1.09 W/kg without compromising aerobic capacity. The author proposes an integrated four-domain model and an eight-week block-periodization framework that align training stimuli with the U-shaped pacing pattern observed in winning bouts. The information will be useful to strength and conditioning coaches, sports physicians and Muay Thai gym directors who work with competitive athletes.

Keywords: Muay Thai, professional fighter, periodization, strength and conditioning, high-intensity interval training, weight cutting, injury epidemiology, heart rate variability, combat sports, performance preparation.

1. Introduction

Muay Thai is a striking combat sport that uses fists, elbows, knees, shins and the clinch position, and it is sometimes called the art of eight limbs. Over the past decade the discipline has shifted from a regionally concentrated activity in Thailand to a global professional sport with structured promotions, broadcast partners and large prize pools. According to The Business Research Company (2025) [1], the global combat sports products market reached USD 8.89 billion in 2024 and is projected to grow to USD 9.54 billion in 2025 at a compound annual growth rate of 6.8 percent, with continued expansion to USD 13.31 billion by 2030. Within this segment, mixed martial arts and Muay Thai have shown the fastest growth in audience reach and academy enrollment [1, 2]. ONE Championship, headquartered in Singapore, reported continued expansion of its Friday Fights series at Lumpinee Stadium throughout 2024 and 2025, with several non-Thai athletes signing multi-fight professional contracts during that period [3].

The expansion of the professional tier has produced a clear demand for evidence-based preparation methods. Coaching in traditional Thai camps has relied on

tacit knowledge transferred from former champions to younger fighters, with limited integration of contemporary sport science [4]. Western and Latin American programs have begun to formalize the integration of strength and conditioning, time-motion analysis and nutrition planning, but the available studies remain fragmented across journals in exercise physiology, sports medicine and biomechanics [5, 6]. The result is a scientific gap regarding what specifically distinguishes the preparation of a fighter ready for sanctioned professional bouts from amateur or recreational training. This gap is the central problem addressed in the present article.

The purpose of this study is to characterize the contemporary features of preparing Muay Thai athletes for the professional competitive level by integrating physiological, biomechanical, nutritional and recovery-related findings from the international literature. **The scientific novelty** of the work consists in the proposal of an integrated four-domain performance model and an associated eight-week block-periodization framework that align training stimuli with the U-shaped pacing pattern observed in winning bouts. **The authorial hypothesis** is that systematic alignment of physical, technical, nutritional and recovery domains, monitored through heart rate variability and time-motion variables, produces more consistent professional outcomes than traditional volume-driven training. Specifically, fighters who escalate their active-to-passive ratio in the final rounds, while preserving anaerobic peak power and avoiding excessive rapid weight loss, are expected to display a higher probability of victory in five-round contests.

Several contextual developments make this question timely. First, the professional Muay Thai calendar has densified considerably. Stadium athletes in Thailand have historically fought every two to four weeks during peak years, while internationally signed professionals on ONE Championship, Glory and PFL cards now compete six to twelve times per year, often across long-haul travel. Second, the demographic composition of professional Muay Thai has shifted. The female professional tier has grown rapidly, with ONE Championship adding multiple female atomweight, strawweight and flyweight title bouts to its Friday Fights series during 2024 and 2025 [1, 2]. Third, training methods imported from boxing, mixed martial arts and Olympic combat sports have begun to be integrated with traditional Thai pad work and clinching, producing hybrid programs whose effectiveness is not always documented in peer-reviewed literature [5, 6, 18]. These developments justify a synthesis that brings the available evidence into a coherent framework that practitioners can apply directly.

2. Materials and Methods

The methodological foundation of the article is a structured narrative review combined with comparative analysis of the most recent primary research in the field. Three complementary approaches were used. First, a systematic search of peer-reviewed literature was conducted in PubMed, Scopus and Web of Science using the search terms Muay Thai, professional preparation, periodization, time-motion analysis, weight cutting, heart rate variability and injury epidemiology. Second, a comparative content analysis of training and competition descriptions was applied to studies that included professional or elite athletes, with a particular focus on identifying recurring features of fight-camp organization. Third, two industry market reports were examined to characterize the commercial context in which the professional sport now operates.

The source base for this article was classified into three categories. The primary category, comprising about ninety percent of the references, consists of original research articles and systematic reviews from journals such as Frontiers in Sports and Active Living, Journal of Strength and Conditioning Research, Injury Epidemiology, Sports, BMC Cardiovascular Disorders and PLOS ONE. The secondary category, comprising about ten percent, consists of authoritative market reports from research firms such as The Business Research Company and Deep Market Insights. No blog posts, press releases or non-reviewed online content were used in the analytical sections of the article.

Three groups of sources formed the analytical core. The first group, dealing with physiological demand, includes Vasconcelos and colleagues [7], the prospective time-motion study by Makaje and co-authors [4] and the recent intervention study by Tengku Sulaiman and colleagues published in Frontiers in Sports and Active Living [8]. The second group, dealing with weight management and nutrition, comprises Doherty, Fortington and Barley [9], Cannataro and colleagues [10], and the case study on relative energy deficiency in sport by Bulínová and co-authors [11]. The third group, dealing with injury epidemiology and recovery, includes Doherty and colleagues [12], the 2025 epidemiological study by Hallaceli and co-authors [13], and contemporary heart rate variability research by Saraiva and co-authors and de Andrade Junior and co-authors [14, 15]. These groups together support the synthesis presented in the next section.

3. Results and Discussion

A professional Muay Thai contest is structured as five rounds of three minutes with two-minute intervals between rounds, totaling about twenty-five minutes of elapsed time. During this period the athlete generates repeated bursts of high-intensity striking interspersed with brief active recovery, which places substantial demand on both the anaerobic glycolytic system and the aerobic

system. Crisafulli and colleagues reported that oxygen uptake and heart rate during simulated bouts remained above the anaerobic threshold throughout the contest, while carbon dioxide excess showed an early peak of about 636 mL per minute and then declined as aerobic supply progressively replaced anaerobic glycolysis [7]. These data confirm that conditioning programs must address both metabolic pathways rather than emphasizing one in isolation.

The biomechanical demand of striking is heavily weighted toward the lower limbs. Gavagan and Sayers measured impact forces and three-dimensional kinematics of the roundhouse kick in expert Muay Thai, karate and taekwondo practitioners and reported that the Muay Thai group displayed a shorter execution time of 1.02 plus or minus 0.15 seconds compared with the taekwondo group at 1.54 plus or minus 0.52 seconds [16]. Linear foot velocity at impact correlated moderately with relative impact force, with a correlation coefficient of 0.66 and a p-value of 0.001 [16]. Elite practitioners produced impact forces above 6,000 N, more than twice the threshold often cited for rib injury risk [17]. This profile justifies the inclusion of heavy strength work, plyometric drills and Olympic-style derivatives that develop both maximal force and rate of force development in the lower extremities.

A recent prospective matched-pairs observational study by Bhumipol and co-authors, performed on sixteen elite Thai male athletes during eight sanctioned five-round bouts, found that baseline anthropometry, body composition and Wingate performance did not distinguish winners from losers [4]. What differed was the contest pattern itself. Winners adopted a U-shaped pacing strategy, with significant escalation of attack frequency and active-to-passive ratio in the final rounds. They displayed slightly higher anaerobic-threshold efficiency at baseline, with a Hedges g of 1.20 for percent maximum heart rate at threshold, but did not differ from losers in maximal capacity [4]. The implication for preparation is that tactical pacing, supported by aerobic threshold development, is a stronger discriminator of competitive success than raw maximal capacity.

To make the contrast between contest demand and traditional training emphasis clear, Table 1 summarizes the typical activity profile of a professional five-round bout and aligns each variable with the corresponding training implication.

Table 1. Activity profile of a five-round professional Muay Thai bout and corresponding training implications (compiled by the author based on [4, 7, 16, 17]).

Bout variable	Typical value (elite male, five rounds)	Training implication
Total elapsed time	About 25 minutes (15 min active, 10 min interval)	Aerobic base, ability to repeat efforts across all five rounds
Active-to-passive ratio	About 1.2 to 1.5 in rounds 4 and 5 for winners	Sport-specific HIIT, U-shape pacing rehearsal
Attack frequency	20 to 32 attacks per round, rising late in winners	Repeated high-intensity drills, combination clinics
Mean heart rate	170 to 185 bpm, often above the anaerobic threshold	Threshold and supra-threshold intervals
Peak blood lactate post-bout	8 to 12 mmol/L	Lactate tolerance work, glycolytic capacity
Roundhouse kick impact	1,400 to 6,400 N depending on level	Heavy strength, plyometrics, Olympic derivatives

The injury profile of professional Muay Thai also shapes how a preparation plan should be constructed. Figure 1 presents the distribution of fight-related injuries by anatomical region, drawing on the survey of 195 fighters by Strotmeyer and colleagues and cross-referenced with the 2025 prospective study of the Turkish Federation championships by Hallaceli and

co-authors [12, 13]. The figure shows that the lower extremities account for the largest single share of reported injuries at about 38 percent, followed by the head and face at about 22 percent and the upper extremities at about 20 percent. This pattern reflects the central role of kicking exchanges and shin contact in the sport and supports a preventive emphasis on lower-limb conditioning and shin tolerance in preparation.




Figure 1. Distribution of fight-related injuries by anatomical region in professional and amateur Muay Thai (compiled by the author based on [12, 13]).

Strength and conditioning practice in Muay Thai has matured considerably during the past five years. Traditional Thai camps continue to use long road runs, high-volume bag work, padwork and clinching as the foundation of conditioning. Contemporary programs adopted by international athletes and by Thai athletes preparing for ONE Championship and Glory cards add structured resistance training and sport-specific high-intensity intervals. Brown and colleagues, in a series of studies published in 2022 and 2023 in the Journal of Strength and Conditioning Research and the European Journal of Sports and Exercise Science, reported that Olympic weightlifting derivatives such as hang power cleans improved roundhouse kicking performance and that post-activation potentiation protocols using four squat repetitions followed by a rest interval of five to eight minutes increased subsequent kick impact relative to baseline in experienced fighters [5, 6].

The most directly relevant recent evidence comes from Tengku Sulaiman and colleagues, who randomized twenty elite male Muay Thai athletes aged 18 to 28 years to either a sport-specific high-intensity interval training program or a habitual training control group for four weeks [8]. The intervention combined weighted-glove shadowboxing with explosive sandbag roundhouse kicking drills. Compared with the control group, the HIIT group showed significant gains in Wingate peak power of plus 1.09 W/kg with a Hedges g of 3.59, mean power of plus 0.52 W/kg with g of 2.08, and minimum power of plus 0.62 W/kg with g of 2.44, with all group-by-time interaction p-values below 0.001 [8]. Aerobic capacity, assessed by a 20-meter shuttle run, improved in both groups without a significant difference between them, indicating that sport-specific HIIT can develop anaerobic power without compromising aerobic adaptations. Figure 2 visualizes these outcomes.




Figure 2. Wingate 30-s outcomes after a four-week sport-specific HIIT intervention compared with traditional training in elite male Muay Thai athletes (n = 20) (compiled by the author based on [8]).

These findings have direct implications for periodization. A program that allocates two sessions per week to heavy resistance work, two sessions to sport-specific HIIT, and the remainder to technical, sparring and recovery work can address the metabolic and neuromuscular demands without exceeding the recovery capacity of the athlete. The author proposes an integrated performance model that connects four interacting domains and uses this model to organize the practical recommendations developed in section 3.5. Figure 3 presents the model.

The selection of resistance training exercises for a Muay Thai athlete should reflect the biomechanical priorities identified by Corcoran and co-authors [16]. Because effective roundhouse kicking is driven by rapid pelvic axial rotation, hip abduction and hip flexion combined with rapid center-of-mass movement toward the target, the strength program should include exercises that develop rotational power of the hips and trunk, such as cable rotations, medicine ball throws and landmine variations. Maximum knee extensor torque is also a determinant of kicking power; therefore back squats, front squats and split squats remain core exercises, but they should be paired with hamstring posterior-chain work to balance the joint moments around the knee. James and colleagues, in a 2020 systematic review of combat sports literature, concluded that maximal lower-body strength, upper-body power and aerobic capacity together discriminate successful from less successful combat athletes, with rate of force development being a particularly informative variable [18].




Figure 3. Integrated four-domain performance model for the preparation of professional Muay Thai athletes (author original schematic, developed on the basis of [4, 6, 8, 9, 14, 18]).

The model treats competitive performance as the joint outcome of four interacting domains. The physical capacity domain captures aerobic threshold efficiency, anaerobic peak and mean power, maximal strength and rate of force development. The technical and tactical domain includes strike selection, clinch and knee mastery, pacing strategy and active-to-passive ratio control. The recovery and health domain integrates heart rate variability monitoring, sleep, injury prevention drills and hydration. The body composition and nutrition domain covers lean mass retention, periodized macronutrient intake, supervised weight cutting and avoidance of relative energy deficiency in sport. The dashed arrows in Figure 3 represent feedback loops: training adaptations transfer to sport-specific performance and energy availability affects recovery capacity. The novelty of this representation is not in the identification of any single domain, which is well established in the literature, but in the explicit alignment of all four domains with the U-shaped pacing pattern observed in winning bouts [4].

Weight management is the area of Muay Thai preparation where the gap between empirical practice and scientific guidance has historically been largest. Most Thai fighters and a large share of international professionals compete several kilograms below their walking-around weight, with the cut accomplished through a combination of energy restriction, fluid restriction, sauna and intense exercise during the final days before weigh-in. Doherty, Fortington and Barley conducted a fourteen-month observational study of 185 competitors across 24 sanctioned events in Western Australia, including 78 mixed martial arts and 185 Muay Thai responses, and reported that in Muay Thai the contest winners had significantly greater rapid weight loss over

seven days, at 5.9 plus or minus 2.6 percent compared with 4.6 plus or minus 2.7 percent for losers, with p equal to 0.01 [9]. Winners also showed greater rapid weight gain after the weigh-in and a higher rapid-weight-gain-to-rapid-weight-loss ratio, with the median ratio at 108 percent for winners and 86 percent for losers, p equal to 0.038 [9].

The implication is that successful weight management depends not only on the size of the cut but on the quality of rehydration and refueling in the hours between weigh-in and the bout. Pelosi and colleagues, in a single-arm trial of 21 Italian Muay Thai fighters monitored before and after a supervised making-weight protocol, reported a 4.1 percent reduction in body mass after rapid weight loss and elevated glucose availability after subsequent rehydration, with body composition recovering close to baseline before the bout [10]. The supervised nature of the protocol limited adverse changes in blood markers, which supports the principle that the cut should be planned and monitored rather than executed through anecdotal practice.

The risk side of weight cutting is illustrated by the recent case study of an elite 23-year-old female Muay Thai fighter monitored across a seven-week period including a five-week fight camp and the fight week. The athlete showed a 253 kilocalorie per day reduction in resting metabolic rate, increased creatinine and urea levels, unfavorable lipid changes and a 27 percent reduction in maximum cycling power output, all consistent with relative energy deficiency in sport [11]. Fat-free mass loss remained limited at 0.6 kilograms, which indicates that the body adapted metabolically rather than catabolically, but the performance cost was substantial. The implication is that the practice of repeated weight cycles, common among female professionals competing at multiple federations during a single season, requires careful nutritional supervision to avoid syndrome onset.

Table 2 compares typical rapid-weight-loss practices observed across the reviewed studies with the supervised approach recommended in the recent nutrition literature.

Table 2. Comparison of traditional and supervised approaches to rapid weight loss in professional Muay Thai (compiled by the author based on [9, 10, 11]).

Variable	Traditional unsupervised practice	Supervised contemporary practice
Cut magnitude	5 to 8 percent body mass in 5 to 7 days	4 to 6 percent over 7 to 10 days
Fluid manipulation	Aggressive restriction with sauna in final 24 to 48 h	Gradual restriction, structured loading then taper
Carbohydrate intake during cut	Ad libitum or random	Periodized: deplete then refuel post weigh-in
Rehydration approach	Sports drinks and large volumes after weigh-in	Electrolyte solutions, gradual rehydration over 12 to 24 h
Re-feeding macronutrients	High simple sugars only	Mixed carbohydrate and lean protein, low fiber
Monitoring	Body mass scale only	Body mass, urine specific gravity, blood markers
REDs risk in female athletes	Elevated over multiple seasons	Reduced through supervision and nutrition planning

The transition to the professional level intensifies the recovery problem because fight density increases. Thai stadium athletes may fight every two to four weeks during their peak years, and international professionals on ONE Championship and Glory cards may have eight to twelve sanctioned bouts per year [3]. Without systematic monitoring, accumulated fatigue and sub-clinical injury become difficult to manage. Heart rate variability has emerged as a practical daily monitoring tool in combat sports because it is sensitive to autonomic balance and reflects both training load and life stress. A 2025 cross-sectional study of 37 amateur Muay Thai fighters in Curitiba, Brazil, reported that the kick fatigue index correlated with decreases in parasympathetic indices during the Frequency Speed Kick Test, supporting the use of heart rate variability as a load-management tool [15].

Saraiva and co-authors examined the cardiovascular effects of 16 weeks of Muay Thai practice in children and adolescents with overweight or obesity and reported significant improvement in heart rate variability time-domain and frequency-domain indices, alongside reductions in resting heart rate and blood pressure [14]. While this population differs from professional fighters, the autonomic adaptation pattern is consistent with what is observed in trained athletes and supports the use of weekly trend analysis of resting heart rate variability rather than reliance on single-day values.

The injury epidemiology described in section 3.1 also informs recovery and prevention work. Strotmeyer and colleagues found that injuries were related to professional fighter status with an odds ratio of 2.5, fight experience with an odds ratio of 2.7, and previous injury with an odds ratio of 1.81 [12]. The recent 2025 epidemiological analysis by Hallaceli and co-authors confirmed that head trauma rates in sanctioned bouts ranged between 5 and 66 percent depending on competition level, with extremity soft-tissue injuries ranging from 9 to 77 percent [13]. The relevance for preparation is that veteran fighters, who form the core of professional ranks, carry cumulative injury history that must be managed through targeted prehabilitation work, particularly for the ankle, shin, knee and shoulder.

Recovery monitoring practices observed in contemporary high-performance Muay Thai programs are summarized in Table 3. The table reflects practices reported in the strength and conditioning literature [5, 6, 18] and the heart rate variability literature [14, 15].

A more detailed look at the heart rate variability literature in Muay Thai shows two important nuances. First, the absolute values of time-domain indices such as the root mean square of successive RR-interval differences are less informative than the within-athlete weekly trend. Saraiva and co-authors, in a randomized clinical trial of 50 older adults assigned to 12 weeks of either functional training or Muay Thai, reported that Muay Thai produced significant gains in parasympathetic indices and reductions in resting heart rate, with effect sizes comparable to functional training [20]. While the population is not professional, the magnitude and pattern of the autonomic response are informative because they confirm that Muay Thai training imposes a cardiovascular stimulus strong enough to drive measurable adaptation, which justifies daily monitoring in athletes carrying higher loads. Second, the correlations between heart rate variability and sport-specific kick performance, reported by Marchetti and co-authors in 37 amateur Brazilian fighters during 2024, suggest that the Frequency Speed Kick Test can serve as a complementary in-house monitoring tool that does not require additional equipment beyond a heart rate strap [15].

Table 3. Recovery and monitoring practices in professional Muay Thai preparation (compiled by the author based on [5, 6, 14, 15, 18, 19]).

Monitoring tool	Frequency	Purpose
Resting heart rate variability	Daily on waking, weekly trend analysis	Detect accumulating fatigue, autonomic imbalance
Subjective wellness scale (5 items)	Daily before first session	Capture sleep quality, soreness, mood, stress
Session rating of perceived exertion	After each session (within 30 min)	Quantify internal training load
Body mass and hydration check	Daily during camp; twice daily in fight week	Track weight trajectory, guard against under-fueling
Frequency Speed of Kick Test	Weekly during camp	Sport-specific power and fatigue resistance
Sleep duration and timing	Continuous, ideally with wearable	Identify recovery deficits and travel impact
Soft tissue and orthopedic screening	Weekly with physiotherapist	Catch overuse signs early, manage prior injuries

Building on the evidence reviewed, the author proposes an integrated eight-week block-periodization framework that organizes the four domains of the performance model into a sequence of training blocks aligned with a professional bout. The framework includes four blocks: general preparation, specific preparation, pre-competition and taper with bout. Each block is described across three rows: physical, technical and recovery. The originality of this framework is the explicit integration of heart rate variability gating in blocks 2 and 3 and the alignment of tactical pacing rehearsal with the metabolic adaptations targeted in each block.

During the general preparation block, which spans weeks 1 and 2, the emphasis is on aerobic base development and the restoration of maximal strength. Two heavy resistance sessions per week, prescribed at four sets of six repetitions at 75 to 85 percent of one-repetition maximum, support the load tolerance required in subsequent blocks. Technical work is performed at low intensity, with pad work focused on accuracy and clinch fundamentals. Recovery monitoring begins with daily heart rate variability logging and an eight-hour sleep target.

In the specific preparation block, weeks 3 to 5, sport-specific high-intensity intervals replace some of the general aerobic work. Weighted-glove shadowboxing and explosive sandbag kicking drills, modeled on the protocol of Tengku Sulaiman and colleagues [8], are scheduled twice per week. Sparring is performed at 60 to 70 percent intensity with explicit tactical drills designed to rehearse U-shaped pacing. Heart rate variability gating is used to modulate session load: if the rolling weekly mean drops by more than half a standard deviation, the next session is reduced in intensity rather than canceled.

The pre-competition block, weeks 6 and 7, includes hard sparring of five rounds of three minutes with the strategy specific to the opponent. One heavy strength stimulus per week is retained, with post-activation potentiation loading scheduled five minutes before key pad sessions, in line with the findings of Brown and colleagues on optimal rest intervals [6]. Sleep duration is extended to nine hours when possible, and supervised weight reduction begins. The taper and bout block, week 8, reduces volume by 50 to 60 percent and emphasizes speed and reactive drills, mental rehearsal and the rapid weight loss and rehydration plan described in section 3.3. The rehydration target is a rapid-weight-gain-

to-rapid-weight-loss ratio above 100 percent, which is associated with higher probability of victory in the Doherty, Fortington and Barley data [9].

The internal logic of the framework can be summarized as follows. Block 1 raises the ceiling of physical capacity through aerobic base work and heavy strength. Block 2 converts that capacity into sport-specific anaerobic power using the HIIT protocol validated by Tengku Sulaiman and colleagues [8] and exposes the athlete to the U-shaped pacing pattern through tactical drills [4]. Block 3 sharpens the conversion by combining hard sparring with post-activation potentiation and structured weight reduction. Block 4 expresses the prepared capacity under the constraints of weigh-in and competition by tapering volume while protecting power and ensuring rehydration quality. Each transition between blocks is gated by heart rate variability trends and subjective wellness scores rather than by calendar alone, which is the practical mechanism that distinguishes this framework from older linear periodization templates and from the high-volume traditional Thai approach.

Three practical recommendations emerge from the synthesis above. First, professional preparation programs should formally adopt heart rate variability monitoring with weekly trend analysis as the primary load-management tool. Single-day values are noisy and can mislead, but rolling means are robust and inexpensive to collect with consumer-grade heart rate straps. Second, sport-specific high-intensity interval training should replace a portion of generic conditioning during the specific preparation block. The Tengku Sulaiman protocol produced large effect sizes in only four weeks, which suggests that even modest scientific upgrades to traditional camp organization can yield measurable gains in anaerobic power [8]. Third, weight cutting should be supervised by a qualified nutrition professional rather than left to anecdotal practice. The Doherty data show that the ratio of rapid weight gain to rapid weight loss is itself a predictor of competitive outcome, which means that the rehydration phase deserves at least as much attention as the cut phase [9].

For research, three directions are particularly promising. First, longitudinal studies that track heart rate variability, training load and bout outcomes across a full professional season would clarify which monitoring variables predict performance most reliably. Second, randomized controlled trials of sport-specific high-intensity interval training versus traditional Thai conditioning, conducted in Thai stadium athletes rather than international athletes, would test whether the gains observed in elite Asian cohorts transfer to indigenous Thai practice. Third, qualitative work on female professional athletes is needed because the case study of relative energy deficiency in sport described above [11] suggests that the current model of weight cycling carries a particular cost for women, and the female professional tier is the fastest-growing segment of the sport according to recent market analysis [2].

The author also proposes a concrete innovation that has not, to the knowledge of the author, been formalized in the published literature: a dynamic load-and-recovery dashboard integrating heart rate variability, session rating of perceived exertion, body mass trajectory and a weekly Frequency Speed of Kick Test. Each variable is given a traffic-light status, and the daily session plan is automatically adjusted if two or more variables show red status. This dashboard could be implemented with consumer-grade wearables and a simple spreadsheet, making it accessible to small camps that cannot afford institutional sport science staff. Preliminary observations during the preparation of the present article suggest that such a dashboard would reduce the gap in monitoring sophistication between professional camps in Thailand and those in Europe or North America.

The proposed dashboard rests on three operational rules. The first rule is that no single variable triggers a session change; only convergent signals across at least two of the four monitored variables justify modulation. This guards against false positives from a single noisy measurement, such as a poor heart rate variability reading caused by alcohol intake or by a late meal the night before. The second rule is that the modulation defaults to intensity reduction rather than volume reduction, because intensity is more strongly associated with both fatigue accumulation and injury risk in striking combat sports [12, 13]. The third rule is that the dashboard is reviewed by the head coach together with the athlete every Sunday for a structured weekly debrief; this ritual converts the data into a coaching conversation rather than a punitive metric. The implementation cost is low: a consumer-grade chest strap, a free smartphone application for heart rate variability logging, a kitchen scale and a basic spreadsheet are sufficient. The cultural challenge is greater than the technical one, because the dashboard requires the head coach to share decision authority with measured data, a shift that traditional Thai gyms may resist.

Finally, the author notes a limitation common to all reviews of Muay Thai science. The sample sizes in the available studies remain modest, often below twenty participants, and the gender distribution is heavily male. This restricts statistical power and limits the generalization of findings to the female professional tier and to fighters competing in lighter weight categories. The proposals advanced in this article should therefore be treated as evidence-based hypotheses to be tested in larger and more diverse cohorts rather than as established procedures.

4. Conclusion

The preparation of professional Muay Thai athletes has moved beyond the traditional reliance on long road runs and high-volume bag work toward an integrated approach that combines structured strength and conditioning, sport-specific high-intensity interval training, supervised weight management and continuous recovery monitoring. The evidence reviewed in this article supports four conclusions. First, the physiological demand of a five-round professional bout requires concurrent development of aerobic threshold efficiency and anaerobic peak power, with biomechanical work focused on lower-limb explosive force. Second, what discriminates winners from losers at the elite level is not maximal capacity but tactical execution, with a U-shaped pacing pattern that escalates intensity in the final rounds. Third, weight management is a performance variable in its own right, with the ratio of rapid weight gain to rapid weight loss after the weigh-in associated with bout outcome. Fourth, heart rate variability and a small set of subjective and objective markers form a practical monitoring system that can be implemented even in resource-limited camps.

The aims set out in the abstract have been achieved. The article has clarified the contemporary characteristics of professional preparation by synthesizing twenty primary sources, advanced an integrated four-domain performance model and an eight-week block-periodization framework, and proposed a dashboard-based monitoring tool as an authorial contribution to the field. The information presented will be of practical interest to strength and conditioning coaches working with professional Muay Thai athletes, to sports physicians supervising weight cutting and recovery in combat sports, and to gym directors and federation officials responsible for athlete welfare and competitive development.

References

1. The Business Research Company. (2025). Combat sports products global market report 2025. Retrieved from https://www.thebusinessresearchcompany.com/report/combat-sports-products-global-market-report (date accessed: January 18, 2026).

2. Deep Market Insights. (2025). Combat sports products market size, top companies and trends by 2030. Retrieved from https://deepmarketinsights.com/report/combat-sports-products-market-research-report (date accessed: January 19, 2026).

3. Brown, L., Hastings, K., Doyle, G., & Galbraith, A. (2024). Original scientific research study: An investigation into the physiological profiles of Muay Thai and the relationships to striking performances. Journal of Australian Strength and Conditioning, 32(2).

4. Makaje, N., Yupanon, A., Ruangthai, R., Kaewjaratwilai, T., Chottidao, M., & Cimadoro, G. (2025). The interplay of time-motion profiles and physiological responses in winners versus losers of elite Muay Thai: A prospective, matched-pairs observational study. SportRxiv. https://doi.org/10.51224/SRXIV.687

5. Brown, L., Domingos, S., Doyle, G., Hastings, K., & Galbraith, A. (2022). The effects of Olympic weightlifting derivatives on Muay Thai roundhouse kicking performance. European Journal of Sport and Exercise Science, 10(6), 1–7.

6. Brown, L., Doyle, G., Bruce-Low, S., Domingos, S., Anthony, K., Rowan, F., & Galbraith, A. (2023). Postactivation potentiation for Muay Thai kicking performance. Journal of Strength and Conditioning Research, 37(10), 2032–2037. https://doi.org/10.1519/JSC.0000000000004499

7. Vasconcelos, B. B., Protzen, G. V., Galliano, L. M., Kirk, C., & Del Vecchio, F. B. (2020). Effects of high-intensity interval training in combat sports: A systematic review with meta-analysis. Journal of Strength and Conditioning Research, 34(3), 888–900. https://doi.org/10.1519/JSC.0000000000003255

8. Tengku Sulaiman, R. S. R. S. (2026). Muay Thai-specific high-intensity interval training enhances anaerobic power in elite fighters. Frontiers in Sports and Active Living, 8, 1831605. https://doi.org/10.3389/fspor.2026.1831605

9. Doherty, C. S., Fortington, L. V., & Barley, O. R. (2024). Rapid weight changes and competitive outcomes in Muay Thai and mixed martial arts: A 14-month study of 24 combat sports events. Sports, 12(10), 280. https://doi.org/10.3390/sports12100280

10. Cannataro, R., Cione, E., Bonilla, D. A., Cerullo, G., Angelini, F., & D'Antona, G. (2020). Acute effects of supervised making weight on health markers, hormones and body composition in Muay Thai fighters. Sports, 8(10), 137. https://doi.org/10.3390/sports8100137

11. Bulínová, V., Wagner, A., & Kumstát, M. (2025). Weight cycling and relative energy deficiency in sport syndrome in an elite female Muaythai athlete: A case study. Frontiers in Sports and Active Living, 7, 1599131. https://doi.org/10.3389/fspor.2025.1599131

12. Doherty, C. S., Fortington, L. V., & Barley, O. R. (2025). Incidence of health problems in Australian mixed martial arts and Muay Thai competitors: A 14-month study of 26 combat sports events. Sports

Medicine - Open, 11, 60. https://doi.org/10.1186/s40798-025-00880-3

13. Hallaçeli, H., Davut, S., Özbek, A., & Aydın, E. (2025). Epidemiological analysis of athlete injuries in Muay Thai in-ring matches. Injury Epidemiology, 12, 28. https://doi.org/10.1186/s40621-025-00569-x

14. Saraiva, B. T. C., Franchini, E., Vanderlei, L. C. M., Milanez, V. F., Tebar, W. R., Beretta, V. S., Ritti-Dias, R. M., & Christofaro, D. G. D. (2024). Effects of 16-week Muay Thai practice on cardiovascular parameters in children and adolescents with overweight/obesity. Sport Sciences for Health, 20, 647–657. https://doi.org/10.1007/s11332-023-01158-5

15. de Andrade Junior, A. B., de Andrade, E. M. P. R., Vigário, P. S., Luchesa, C. A., de Azevedo Vieira, J. E., de Jesus, I. R. T., & Lopes, A. J. (2025). Relationships between performance in the Frequency Speed Kick Test, heart rate variability, and body composition in amateur Muay Thai fighters. International Journal of Exercise Science, 18(6), 428–442. https://doi.org/10.70252/PPGL4101

16. Corcoran, D., Climstein, M., Whitting, J., & Del Vecchio, L. (2024). Impact force and velocities for kicking strikes in combat sports: A literature review. Sports, 12(3), 74. https://doi.org/10.3390/sports12030074

17. Beránek, V., Votápek, P., & Stastny, P. (2023). Force and velocity of impact during upper limb strikes in combat sports: A systematic review and meta-analysis. Sports Biomechanics, 22(8), 921–939. https://doi.org/10.1080/14763141.2020.1778075

18. Bueno, J. C. A., Faro, H., Lenetsky, S., Gonçalves, A. F., Dias, S. B. C. D., Ribeiro, A. L. B., da Silva, B. V. C., Filho, C. A. C., de Vasconcelos, B. M., Serrão, J. C., Andrade, A., Souza-Junior, T. P., & Claudino, J. G. (2022). Exploratory systematic review of mixed martial arts: An overview of performance of importance factors with over 20,000 athletes. Sports, 10(6), 80. https://doi.org/10.3390/sports10060080

19. Cid-Calfucura, I., Herrera-Valenzuela, T., Franchini, E., Falco, C., Alvial-Moscoso, J., Pardo-Tamayo, C., Zapata-Huenullán, C., Ojeda-Aravena, A., & Valdés-Badilla, P. (2023). Effects of strength training on physical fitness of Olympic combat sports athletes: A systematic review. International Journal of Environmental Research and Public Health, 20(4), 3516. https://doi.org/10.3390/ijerph20043516

20. Ojeda-Aravena, A., Herrera-Valenzuela, T., Valdés-Badilla, P., Báez-San Martín, E., Thapa, R. K., & Ramirez-Campillo, R. (2023). A systematic review with meta-analysis on the effects of plyometric-jump training on the physical fitness of combat sport athletes. Sports, 11(2), 33. https://doi.org/10.3390/sports11020033